


PENDING CLAIMS WITH ENTRY OF THE AMENDMENT

1. A compound of the formula:



wherein:


R<sup>1</sup> is a member selected from the group consisting of hydrogen and optionally substituted C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sup>2</sup> is a member selected from the group consisting of amino and mono- or di-substituted amino wherein the substituent is a protecting group;

R<sup>3</sup> is optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>4</sup> is a member selected from the group consisting of hydrogen and L;

L is of the formula



wherein:

R<sup>5</sup> is a member selected from the group consisting of hydrogen, hydroxyl, and substituted hydroxyl wherein the substituent is a protecting group;

R<sup>6</sup> is a member selected from the group consisting of hydrogen, phosphoramidite, an H-phosphonate, a methyl phosphonate, a phosphorothioate, a phosphotriester, a hemisuccinate, a hemisuccinate covalently bound to a solid support, a dicyclohexylcarbodiimide, and a dicyclohexylcarbodiimide covalently bound to a solid support, a hydroxyalkyl, and a hydroxyalkyl covalently bound to a solid support; and

R<sup>7</sup> is a member selected from the group consisting of hydrogen, a phosphate, a triphosphate, and a protecting group;

with the proviso that R<sup>1</sup> and R<sup>4</sup> are not simultaneously L.

09786666, 061501

a2

Sub  
B1

2. A compound in accordance with claim 1, wherein R<sup>1</sup> is hydrogen; R<sup>2</sup> is a member selected from the group consisting of amino, mono-, and di-substituted amino wherein the substituents are members selected from the group consisting of benzoyl, isobutyryl, phthaloyl, di-n-butylaminomethylidene, dimethylaminomethylidene, p-nitrophenylethoxycarbonyl and dimethylaminomethylenamino;

R<sup>4</sup> is L;

R<sup>5</sup> is a member selected from the group consisting of hydrogen, hydroxyl, hydroxyl substituted with a member selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl, tetrahydropyran-1-yl, 4-methoxytetrahydropyran-4-yl, 1-(2-chloro-4-methyl)phenyl-4-methoxypiperidin-4-yl, t-butyldimethylsilyl, p-nitrophenylethylsulfonyl, tetrahydropyranyl, 4-methoxytetrahydropyranyl, 2-nitrobenzyl, 9-phenylxanthen-9-yl and p-nitrophenylethyl;

R<sup>6</sup> is a member selected from the group consisting of consisting of hydrogen, phosphoramidite, H-phosphonate, hemisuccinate, and hemisuccinate covalently bound to a solid support; and

R<sup>7</sup> is a member selected from the group consisting of hydrogen, trityl, monomethoxytrityl, dimethoxytrityl, phthaloyl, di-n-butylaminomethylene, dimethylaminomethylidene and triphosphate.

3. A compound in accordance with claim 2, wherein R<sup>2</sup> is a member selected from the group consisting of amino and an amino group mono-substituted by a protecting group selected from the group consisting of di-n-butylaminomethylidene, p-nitrophenylethoxycarbonyl, and dimethylaminomethylenamino;

R<sup>5</sup> is a member selected from the group consisting of hydrogen, hydroxyl and hydroxyl substituted with a member selected from the group consisting of dimethoxytrityl, tetrahydropyran-1-yl, t-butyldimethylsilyl, 2-nitrobenzyl, and p-nitrophenylethylsulfonyl;

R<sup>6</sup> is a member selected from the group consisting of hydrogen,  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite and a hemisuccinate covalently bound to controlled pore glass; and

R<sup>7</sup> is a member selected from the group consisting of dimethoxytrityl, di-n-butylaminomethylene, and dimethylaminomethylidene.

09780066-061501

cont'd

a2

4. A compound in accordance with claim 2, wherein R<sup>2</sup> is a member selected from the group consisting of amino and an amino group mono-substituted by a protecting group selected from the group consisting of di-n-butylaminomethylidene, p-nitrophenylethoxycarbonyl, and dimethylaminomethylenamino;

R<sup>5</sup> is a member selected from the group consisting of hydrogen and hydroxyl substituted with a member selected from the group consisting of dimethoxytrityl, tetrahydropyran-1-yl, t-butyldimethylsilyl, 2-nitrobenzyl, and p-nitrophenylethyl;

R<sup>6</sup> is a member selected from the group consisting of hydrogen and  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite; and

R<sup>7</sup> is a member selected from the group consisting of hydrogen and dimethoxytrityl.

5. A compound in accordance with claim 2, wherein R<sup>2</sup> is a member selected from the group consisting of amino and dimethylaminomethylenamino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite; and

R<sup>7</sup> is dimethoxytrityl.

6. A compound in accordance with claim 2, wherein R<sup>2</sup> is amino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite; and

R<sup>7</sup> is dimethoxytrityl.

7. A compound in accordance with claim 2, wherein R<sup>2</sup> is amino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is hydrogen; and

R<sup>7</sup> is hydrogen.

09786666-061501

cont'd

8. A compound in accordance claim 2, wherein R<sup>2</sup> is dimethylaminomethylenamino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite; and

R<sup>7</sup> is dimethoxytrityl.

9. A compound in accordance with claim 2, wherein R<sup>2</sup> is amino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite; and

R<sup>7</sup> is a triphosphate.

10. A compound in accordance with claim 1, wherein;

R<sup>1</sup> is optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>2</sup> is a member selected from the group consisting of amino, mono-, and di-substituted amino wherein the substituent is a member selected from the group consisting of benzoyl, isobutyryl, phthaloyl, di-n-butylaminomethylidene, dimethylaminomethylidene, p-nitrophenylethoxycarbonyl and dimethylaminomethylenamino;

R<sup>3</sup> is optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>4</sup> is L;

R<sup>5</sup> is a member selected from the group consisting of hydrogen, hydroxyl and hydroxyl substituted with a member selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl, tetrahydropyran-1-yl, 4-methoxytetrahydropyran-4-yl, 1-(2-chloro-4-methyl)phenyl-4-methoxypiperidin-4-yl, t-butyldimethylsilyl, p-nitrophenylethylsulfonyl, tetrahydropyranyl, 4-methoxytetrahydropyranyl, 2-nitrobenzyl, 9-phenylxanthen-9-yl and p-nitrophenylethyl;

R<sup>6</sup> is a member selected from the group consisting of hydrogen, H-phosphonate, phosphoramidite, hemisuccinate, and hemisuccinate covalently bound to a solid support; and

09786666-061501

Cont'd

B.  
B2

*a2*

$R^7$  is a member selected from the group consisting of hydrogen, trityl, monomethoxytrityl, dimethoxytrityl, phthaloyl, di-n-butylaminomethylene, and dimethylaminomethylidene.

11. A compound in accordance with claim 10 wherein  $R^1$  is methyl;  $R^2$  is a member selected from the group consisting of amino and an amino group mono-substituted by a protecting group selected from the group consisting of di-n-butylaminomethylidene, p-nitrophenylethoxycarbonyl, and dimethylaminomethylenamino;

$R^3$  is methyl;

$R^5$  is a member selected from the group consisting of hydrogen, hydroxyl and hydroxyl substituted with a member selected from the group consisting of dimethoxytrityl, tetrahydropyran-1-yl, t-butyldimethylsilyl, 2-nitrobenzyl, and p-nitrophenylethylsulfonyl;

$R^6$  is a member selected from the group consisting of hydrogen,  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite and a hemisuccinate covalently bound to controlled pore glass; and

$R^7$  is a member selected from the group consisting of dimethoxytrityl, di-n-butylaminomethylene, and dimethylaminomethylidene.

*cont'd*

12. A compound in accordance claim 10, wherein  $R^1$  is methyl;  $R^2$  is a member selected from the group consisting of amino and an amino group mono-substituted by a protecting group selected from the group consisting of di-n-butylaminomethylidene, p-nitrophenylethoxycarbonyl, and dimethylaminomethylenamino;

$R^5$  is a member selected from the group consisting of hydrogen and hydroxyl substituted with a member selected from the group consisting of dimethoxytrityl, tetrahydropyran-1-yl, t-butyldimethylsilyl, 2-nitrobenzyl, and p-nitrophenylethylsulfonyl;

$R^6$  is a member selected from the group consisting of consisting of hydrogen and  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite; and

$R^7$  is a member selected from the group consisting of hydrogen and dimethoxytrityl.

13. A compound in accordance with claim 10, wherein R<sup>1</sup> is methyl;  
R<sup>2</sup> is a member selected from the group consisting of amino and  
dimethylaminomethylenamino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite; and

R<sup>7</sup> is dimethoxytrityl.

14. A compound in accordance with claim 10, wherein R<sup>1</sup> is methyl;  
R<sup>2</sup> is amino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite; and

R<sup>7</sup> is dimethoxytrityl.

15. A compound in accordance with claim 10, wherein R<sup>1</sup> is methyl;  
R<sup>2</sup> is amino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is hydrogen; and

R<sup>7</sup> is hydrogen.

16. A compound in accordance with claim 10, wherein R<sup>1</sup> is methyl; is  
dimethylaminomethylenamino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite; and

R<sup>7</sup> is dimethoxytrityl.

17. A compound in accordance with claim 10, wherein R<sup>1</sup> is methyl;  
R<sup>2</sup> is amino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen;


R<sup>6</sup> is  $\beta$ -cyanoethyl-N-diisopropyl phosphoramidite; and

09786666 .061501

Q2  
Cont'd

*RP*  
R<sup>7</sup> is a triphosphate.

*SUB B3*  
18. An oligonucleotide comprising one or more nucleotide monomers,  
said monomers having the formula



wherein:


R<sup>1</sup> is a member selected from the group consisting of hydrogen and optionally substituted C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sup>2</sup> is a member selected from the group consisting of amino and mono- or di-substituted amino wherein the substituent is a protecting group;

R<sup>3</sup> is optional substituted C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>4</sup> is L;

L is of the formula



wherein:

R<sup>5</sup> is a member selected from the group consisting of hydrogen and hydroxyl;

R<sup>6</sup> is a member selected from the group consisting of hydrogen, a phosphate, a phosphate covalently attached to a nucleotide, a phosphate covalently attached to a nucleoside; a hemisuccinate covalently bound to a solid support, a dicyclohexylcarbodiimide covalently bound to a solid support, and a hydroxyalkyl covalently bound to a solid support; and

R<sup>7</sup> is a member selected from the group consisting of hydrogen, a phosphate, a phosphate covalently attached to a nucleotide and a phosphate covalently attached to a nucleoside;

09786666-061501

*Contd*

with the proviso that R<sup>1</sup> and R<sup>4</sup> are not simultaneously L.

19. An oligonucleotide in accordance with claim 18, wherein:

R<sup>1</sup> is hydrogen;

R<sup>2</sup> is amino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen and hydroxyl;

R<sup>6</sup> is hydrogen; and

R<sup>7</sup> is a phosphate.

20. An oligonucleotide in accordance with claim 19, wherein:

R<sup>5</sup> is hydrogen.

21. An oligonucleotide in accordance with claim 19 wherein:

R<sup>5</sup> is hydroxyl.

22. An oligonucleotide in accordance with claim 18, wherein:

R<sup>1</sup> is optionally substituted C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sup>2</sup> is amino;

R<sup>3</sup> is methyl;

R<sup>5</sup> is hydrogen and hydroxyl;

R<sup>6</sup> is hydrogen; and

R<sup>7</sup> is a phosphate.

23. An oligonucleotide in accordance with claim 22, wherein

R<sup>1</sup> is methyl and

R<sup>5</sup> is hydrogen.

24. An oligonucleotide in accordance with claim 22, wherein

R<sup>1</sup> is methyl and

R<sup>5</sup> is hydroxyl.

25. An oligonucleotide in accordance with claim 18, wherein said

nucleotide monomers are at the 3' end of said oligonucleotide.

09786666-061501

SUB  
B4

SUB  
C2




26. An oligonucleotide in accordance with claim 18, wherein said nucleotide monomers are at the 5' end of said oligonucleotide.

27. An oligonucleotide in accordance with claim 18, wherein said nucleotide monomers are surrounded by 1 to 10 pyrimidine monomers.

28. An oligonucleotide in accordance with claim 18, wherein said oligonucleotide is a member selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21 and SEQ ID NO:22.

29. A method of detecting the presence, absence, or quantity of a target nucleic acid, said method comprising the steps of:

a) contacting said target nucleic acid with a nucleic acid probe wherein said nucleic acid probe comprises compound of the formula:



wherein:


R<sup>1</sup> is a member selected from the group consisting of hydrogen and optionally substituted C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sup>2</sup> is a member selected from the group consisting of amino and mono- or di-substituted amino wherein the substituent is a protecting group;

R<sup>3</sup> is optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl;

R<sup>4</sup> is L;

L is of the formula



wherein:

$R^5$  is a member selected from the group consisting of hydrogen and hydroxyl;

$R^6$  is a member selected from the group consisting of hydrogen, phosphoramidite, an H-phosphonate, a methyl phosphonate, a phosphorothioate, a phosphotriester, a hemisuccinate, a hemisuccinate covalently bound to a solid support, a dicyclohexylcarbodiimide, and a dicyclohexylcarbodiimide covalently bound to a solid support; and

$R^7$  is phosphate;

with the proviso that  $R^1$  and  $R^4$  are not simultaneously L; located in said probe such that, when said probe hybridizes to said target nucleic acid said compound is in a loop that does not participate in complementary base pairing with a nucleotide of said target nucleic acid; and

b) detecting the fluorescence produced by said fluorescent nucleotide when said probe forms a hybrid duplex with said target nucleic acid.

30. A method of claim 29, wherein said loop ranges in length from about 1 to about 100 nucleotides when said probe hybridizes to said target nucleic acid.

31. A method of claim 29, wherein said loop is an insertion in said nucleic acid probe which is otherwise complementary to said target nucleic acid or to a contiguous subsequence of said target nucleic acid.

32. A method of claim 31, wherein said insertion is three nucleotides in length and comprises two nucleotides each adjacent to said compound.

33. A method of claim 32, wherein at least one nucleotide adjacent to said compound is a purine.

34. A method of claim 33, wherein at least one nucleotide adjacent to said compound is an adenosine.

35. A method of claim 32, wherein at least one nucleotide adjacent to said compound is a pyrimidine.

A2

09780666-064501

Cont'd

36. A method of claim 35, wherein at least one nucleotide adjacent to said compound is a cytosine.

37. A method of claim 34, wherein said compound is bordered by at least two adjacent purines in both the 5' and 3' direction.

38. A method of claim 37, wherein said adjacent purines are adenosine.

39. A method of claim 31, wherein said insertion is said compound.

40. A method of claim 31, wherein said insertion is self-complementary and forms a hairpin wherein said compound is present in the loop of said hairpin and does not participate in complementary base pairing.

41. A method of claim 29, wherein the nucleotides comprising said loop are selected such that they are not complementary to the corresponding nucleotides of the target nucleic acid when said probe is hybridized to said target nucleic acid and wherein said probe is complementary to at least two non-contiguous subsequences of said target nucleic acid.

42. A method of claim 29, wherein said fluorescent nucleotide is present in a terminal subsequence of said nucleic acid probe wherein said terminal subsequence does not hybridize to said target nucleic acid when the remainder of said nucleic acid probe hybridizes to said target nucleic acid.

43. A method of claim 42, wherein said terminal subsequence forms a terminal hairpin by hybridization with a second subsequence of said probe such that said fluorescent nucleotide is present in a loop of said hairpin and does not participate in complementary base pairing.

44. A method of claim 29, wherein said detecting comprises detecting an increase in fluorescence of said fluorescent nucleotide when said probe forms a hybrid duplex with said target nucleic acid.

Q2  
09786666, 061501  
Cont. 20

A2.1  
concl

45. A kit for the detection of nucleic acid-nucleic acid interactions comprising a container, said container containing a compound in accordance with claim 1, and instructions for use.

SF 1233603 v1

09786666 .061501